


Free Green's function - cylindrical geometry

① In preparation towards finding Green's function for



$\vec{r} = (\rho, \phi, z)$

we consider

$-\vec{\nabla} \cdot \epsilon(\vec{r}) \vec{\nabla} G(\vec{r}, \vec{r}') = \delta^{(3)}(\vec{r} - \vec{r}')$

② Let  $\epsilon(\vec{r}) = \epsilon_0$

$-\epsilon_0 \nabla^2 G(\vec{r}, \vec{r}') = \delta^{(3)}(\vec{r} - \vec{r}')$

③  $G(\vec{r}, \vec{r}') = \int_{-\infty}^{+\infty} \frac{dk_z}{2\pi} e^{ik_z(z-z')} \frac{1}{2\pi} \sum_{m=-\infty}^{+\infty} e^{im(\phi-\phi')} g_m(\rho, \rho'; k_z)$

$\delta^{(3)}(\vec{r} - \vec{r}') = \delta(z-z') \delta(\phi-\phi') \frac{\delta(\rho-\rho')}{\rho}$   
 $= \int_{-\infty}^{+\infty} \frac{dk_z}{2\pi} e^{ik_z(z-z')} \frac{1}{2\pi} \sum_{m=-\infty}^{+\infty} e^{im(\phi-\phi')} \frac{\delta(\rho-\rho')}{\rho}$

$\nabla^2 = \frac{1}{\rho} \frac{\partial}{\partial \rho} \rho \frac{\partial}{\partial \rho} + \frac{1}{\rho^2} \frac{\partial^2}{\partial \phi^2} + \frac{\partial^2}{\partial z^2}$

④ Thus, we have

$$- \epsilon_0 \left[ \frac{1}{\rho} \frac{\partial}{\partial \rho} \rho \frac{\partial}{\partial \rho} + \frac{(im)^2}{\rho^2} + (ik_z)^2 \right] g_m(\rho, \rho'; k_z) = \frac{\delta(\rho - \rho')}{\rho}$$

$$\left[ - \frac{1}{\rho} \frac{\partial}{\partial \rho} \rho \frac{\partial}{\partial \rho} + \frac{m^2}{\rho^2} + k_z^2 \right] g_m(\rho, \rho'; k_z) = \frac{1}{\epsilon_0} \frac{\delta(\rho - \rho')}{\rho}$$


⑤ For  $\rho = \rho'$  we can write

$$g_m(\rho, \rho'; k_z) = \begin{cases} A I_m(k_z \rho) + B K_m(k_z \rho) & 0 \leq \rho < \rho' \\ C I_m(k_z \rho) + D K_m(k_z \rho) & 0 \leq \rho' < \rho \end{cases}$$

⑥ Requiring  $g_m(\pm \infty, \rho'; k_z) = 0$  we immediately have

$B = 0$  and  $C = 0$ .

$I_m \rightarrow e^{kz}$   
 $K_m \rightarrow e^{-kz}$



⑦ Integrating around  $\rho = \rho'$  we obtain the boundary conditions

$$- \rho \frac{\partial}{\partial \rho} g_m(\rho, \rho'; k_z) \Big|_{\rho = \rho' - \delta}^{\rho = \rho' + \delta} = \frac{1}{\epsilon_0}$$

and

$$g_m(\rho, \rho'; k_z) \Big|_{\rho = \rho' - \delta}^{\rho = \rho' + \delta} = 0$$

⑧ Using ⑤, ⑥ and ⑦ we have.

$$D K_m(k_z s') - A I_m(k_z s') = 0$$

$$D \frac{d}{ds'} K_m(k_z s') - A \frac{d}{ds'} I_m(k_z s') = -\frac{1}{\epsilon_0 s'}$$

$$⑨ \quad A = -\frac{1}{\epsilon_0 s'} \frac{K_m(k_z s')}{\left[ I_m(k_z s') \frac{d}{ds'} K_m(k_z s') - K_m(k_z s') \frac{d}{ds'} I_m(k_z s') \right]}$$

$$D = -\frac{1}{\epsilon_0 s'} \frac{I_m(k_z s')}{\left[ I_m(k_z s') \frac{d}{ds'} K_m(k_z s') - K_m(k_z s') \frac{d}{ds'} I_m(k_z s') \right]}$$

⑩ We notice the appearance of the Wronskian.

$$K_m \times \left[ -\frac{1}{s} \frac{\partial}{\partial s} \circ \frac{\partial}{\partial s} + \frac{m^2}{s^2} + k_z^2 \right] I_m(k_z s) = 0$$

$$I_m \times \left[ -\frac{1}{s} \frac{\partial}{\partial s} \circ \frac{\partial}{\partial s} + \frac{m^2}{s^2} + k_z^2 \right] K_m(k_z s) = 0$$

Thus,

$$I_m(k_z s) \frac{1}{s} \frac{d}{ds} \circ \frac{d}{ds} K_m(k_z s) - K_m(k_z s) \frac{1}{s} \frac{d}{ds} \circ \frac{d}{ds} I_m(k_z s) = 0$$

(11) Let  $k_2 s = t$

$$I_m \frac{d}{dt} t \frac{d}{dt} K_m - K_m \frac{d}{dt} t \frac{d}{dt} I_m = 0$$

$$I_m \frac{d}{dt} (t K_m') - K_m \frac{d}{dt} (t I_m') = 0$$

$$\frac{d}{dt} [I_m t K_m' - K_m t I_m'] = 0$$

$$\Rightarrow I_m K_m' - K_m I_m' = \frac{c}{t}$$

$c$  - constant.

(12) For  $t \gg 1$  we have.

$$I_m(t) \sim \frac{1}{\sqrt{2\pi}} \frac{1}{\sqrt{t}} e^t$$

$$K_m(t) \sim \sqrt{\frac{\pi}{2}} \frac{1}{\sqrt{t}} e^{-t}$$

$$\begin{aligned} I_m K_m' - K_m I_m' &= \frac{1}{2} \frac{1}{\sqrt{t}} e^t \frac{d}{dt} \left( \frac{1}{\sqrt{t}} e^{-t} \right) - \frac{1}{2} \frac{1}{\sqrt{t}} e^{-t} \frac{d}{dt} \left( \frac{1}{\sqrt{t}} e^t \right) \\ &= -\frac{1}{t} \end{aligned}$$

$$\Rightarrow c = -1$$

Thus,

$$I_m K_m' - K_m I_m' = -\frac{1}{t}$$

(13) Using (12) in (9)

$$A = \frac{1}{\epsilon_0} K_m(k_2 \rho')$$

$$D = \frac{1}{\epsilon_0} I_m(k_2 \rho')$$

(14) Using (13) in (5)

$$g_m(\rho, \rho'; k_2) = \begin{cases} \frac{1}{\epsilon_0} K_m(k_2 \rho') I_m(k_2 \rho) & 0 \leq \rho < \rho' \\ \frac{1}{\epsilon_0} I_m(k_2 \rho') K_m(k_2 \rho) & 0 \leq \rho' < \rho \end{cases}$$

$$= \frac{1}{\epsilon_0} I_m(k_2 \rho_<) K_m(k_2 \rho_>)$$

where

$$\rho_< = \text{Minimum}(\rho, \rho')$$

$$\rho_> = \text{Maximum}(\rho, \rho')$$