

Homework No. 02 (2024 Spring)

PHYS 510: CLASSICAL MECHANICS

School of Physics and Applied Physics, Southern Illinois University–Carbondale

Due date: Tuesday, 2024 Feb 6, 4.30pm

1. (20 points.) The principal identity of functional differentiation is

$$\frac{\delta u(x)}{\delta u(x')} = \delta(x - x'), \quad (1)$$

which states that the variation in the function u at x is independent of the variation in the function u at x' unless $x = x'$. This is a generalization of the identity in multivariable calculus

$$\frac{\partial u^j}{\partial u^i} = \delta_i^j, \quad (2)$$

which states that the variables u^i and u^j are independent unless $i = j$. Using the property of δ -function,

$$\int_{-\infty}^{\infty} dx a(x) \delta(x - x') = a(x'), \quad (3)$$

derive the following identities by repeatedly differentiating by parts.

(a)

$$\int_{-\infty}^{\infty} dx a(x) \frac{d}{dx} \delta(x - x') = -\frac{d}{dx'} a(x') \quad (4)$$

(b)

$$\int_{-\infty}^{\infty} dx a(x) \frac{d^2}{dx^2} \delta(x - x') = +\frac{d^2}{dx'^2} a(x') \quad (5)$$

(c)

$$\int_{-\infty}^{\infty} dx a(x) \frac{d^3}{dx^3} \delta(x - x') = -\frac{d^3}{dx'^3} a(x') \quad (6)$$

(d)

$$\int_{-\infty}^{\infty} dx a(x) \frac{d^n}{dx^n} \delta(x - x') = (-1)^n \frac{d^n}{dx'^n} a(x') \quad (7)$$

2. (20 points.) Evaluate the functional derivative

$$\frac{\delta F[u]}{\delta u(x)} \quad (8)$$

of the following functionals, assuming no variation at the end points.

(a)

$$F[u] = \int_{x_1}^{x_2} dx a(x)u(x) \quad (9)$$

(b)

$$F[u] = \int_{x_1}^{x_2} dx a(x)u(x)^2 \quad (10)$$

(c)

$$F[u] = \int_{x_1}^{x_2} dx \sqrt{1 + u(x)^2} \quad (11)$$

(d)

$$F[u] = \int_{x_1}^{x_2} dx [u(x) + a(x)] [u(x) + b(x)] \quad (12)$$

(e)

$$F[u] = \int_{x_1}^{x_2} dx \frac{a(x)u(x)}{[1 + b(x)u(x)]} \quad (13)$$

3. **(20 points.)** [Refer: Gelfand and Fomin, Calculus of Variations.] Evaluate the functional derivative

$$\frac{\delta F[y]}{\delta y(x)} \quad (14)$$

of the following functionals, assuming no variation at the end points.

(a)

$$F[y] = \int_0^1 dx \frac{dy}{dx} \quad (15)$$

(b)

$$F[y] = \int_{x_1}^{x_2} dx a(x) \frac{dy(x)}{dx} \quad (16)$$

(c)

$$F[y] = \int_0^1 dx y \frac{dy}{dx} \quad (17)$$

(d)

$$F[y] = \int_0^1 dx xy \frac{dy}{dx} \quad (18)$$

(e)

$$F[y] = \int_a^b \frac{dx}{x^3} \left(\frac{dy}{dx} \right)^2 \quad (19)$$

4. (20 points.) Evaluate the functional derivative

$$\frac{\delta F[u]}{\delta u(x)} \quad (20)$$

of the following functionals, assuming no variation at the end points. Given $a(x)$ is a known function.

(a)

$$F[u] = \int_{x_1}^{x_2} dx a(x) \left[1 + \frac{du(x)}{dx} + \frac{d^2u(x)}{dx^2} + \frac{d^3u(x)}{dx^3} \right] \quad (21)$$

(b)

$$F[u] = \int_a^b dx \frac{1}{\left(1 + \frac{d^3u}{dx^3}\right)} \quad (22)$$

(c)

$$F[u] = \int_a^b dx x^5 \sqrt{1 + \frac{d^3u}{dx^3}} \quad (23)$$

(d)

$$F[u] = \int_a^b dx \sqrt{1 + \frac{du}{dx} + \frac{d^3u}{dx^3}} \quad (24)$$

5. (20 points.) Evaluate the functional derivative

$$\frac{\delta W[u]}{\delta u(t)} \quad (25)$$

of the following functionals, with u replaced with the appropriate variable, assuming no variation at the end points.

(a) Let $x(t)$ be position at time t of mass m . The action

$$W[x] = \int_{t_1}^{t_2} dt \frac{1}{2} m \left(\frac{dx}{dt} \right)^2 \quad (26)$$

is a functional of position.

(b) Let $z(t)$ be the vertical height at time t of mass m in a uniform gravitational field g . The action

$$W[z] = \int_{t_1}^{t_2} dt \left[\frac{1}{2} m \left(\frac{dz}{dt} \right)^2 - mgz \right] \quad (27)$$

is a functional of the vertical height.

- (c) Let $x(t)$ be the stretch at time t of a spring of spring constant k attached to a mass m . The action

$$W[x] = \int_{t_1}^{t_2} dt \left[\frac{1}{2}m \left(\frac{dx}{dt} \right)^2 - \frac{1}{2}kx^2 \right] \quad (28)$$

is a functional of the stretch.

- (d) Let $r(t)$ be the radial distance at time t of mass m released from rest in a gravitational field of a planet of mass M . The action

$$W[r] = \int_{t_1}^{t_2} dt \left[\frac{1}{2}m \left(\frac{dr}{dt} \right)^2 + \frac{GMm}{r} \right] \quad (29)$$

is a functional of the radial distance.

- (e) Let $r(t)$ be the radial distance at time t of charge q_1 of mass m released from rest in an electrostatic field of another charge of charge q_2 . The action

$$W[r] = \int_{t_1}^{t_2} dt \left[\frac{1}{2}m \left(\frac{dr}{dt} \right)^2 - \frac{1}{4\pi\epsilon_0} \frac{q_1q_2}{r} \right] \quad (30)$$

is a functional of the radial distance.